

Math Review Summary

CSc 245, Summer 2020

This is a summary of important math concepts from the math review appendix from Dr. McCann's book. For a more detailed review, please read the appendix (on the course webpage).

1 Fractions

Common Fraction Equalities

$$(a) \frac{x}{z} + \frac{y}{z} = \frac{x+y}{z} \quad (b) \frac{x}{z} - \frac{y}{z} = \frac{x-y}{z} \quad (c) \frac{x}{z} \frac{y}{z} = \frac{xy}{z^2} \quad (d) \frac{\frac{x}{z}}{\frac{y}{z}} = \frac{x}{y}$$


$$(e) \frac{x}{w} + \frac{y}{z} = \frac{xz+yw}{wz} \quad (f) \frac{x}{w} - \frac{y}{z} = \frac{xz-yw}{wz} \quad (g) \frac{x}{w} \frac{y}{z} = \frac{xy}{wz} \quad (h) \frac{\frac{x}{w}}{\frac{y}{z}} = \frac{xz}{wy}$$

2 Rational Numbers

Rational Number: A value that can be expressed as the ratio of two integers

3 Set Basics

- **Set**: an unordered collection of unique objects $S = \{x_1, x_2, \dots\}$
- **Notation**:
 - $s \in S$ s is a member of S
 - \emptyset is the empty set ($S = \{\}$)
 - $\{ \text{variables} \mid \text{constraints for membership} \}$ ("variables such that they satisfy the constraints for membership")
 - \mathcal{U} is the universal set (all objects that could possibly be in the set)
- **Operators**:
 - **Union**: $A \cup B$, all objects in A or B (or both)
 - **Intersection**: $A \cap B$, all objects in both A and B
 - **Difference**: $A - B$, all objects in both A that are not also in B
 - **Complement**: \bar{A} , all objects in \mathcal{U} that are not in A ($\mathcal{U} - A$)
 - **Cardinality**: $|A|$, the number of objects in A
- **Venn Diagram**:



- **Notations of Sets of Numbers:**

- \mathbb{Z} : All integers $\{\dots, -2, -1, 0, 1, 2, \dots\}$
- \mathbb{Z}^+ : Positive integers $\{1, 2, 3, \dots\}$
- \mathbb{Z}^* : Non-negative integers $\{0, 1, 2, 3, \dots\}$
- \mathbb{Z}^- : Negative integers $\{\dots, -3, -2, -1\}$
- \mathbb{Z}^{even} : Even integers $\{\dots, -4, -2, 0, 2, 4, \dots\}$
- \mathbb{Z}^{odd} : Odd integers $\{\dots, -3, -1, 1, 3, \dots\}$
- \mathbb{Q} : Rational numbers
- $\overline{\mathbb{Q}}$: Irrational numbers
- \mathbb{R} : all real numbers

4 Associative, Commutative, Distributive, and Transitive properties

- **Associative:** An operation \diamond is associative if $a \diamond (b \diamond c) = (a \diamond b) \diamond c$
- **Commutative:** An operation \diamond is commutative if $a \diamond b = b \diamond a$
- **Distributive:** Operations \diamond and \square are distributive if:
$$a \square (b \diamond c) = (a \square b) \diamond (a \square c)$$
 (\square is left-distributive over \diamond) and
$$(b \diamond c) \square a = (b \square a) \diamond (c \square a)$$
 (\square is right-distributive over \diamond)
- **Transitive:** An relationship \circ is transitive if whenever $a \circ b$ and $b \circ c$, then $a \circ c$ (e.g. $a < b$ and $b < c$ implies $a < c$).

5 Properties of Inequalities

- **Addition:** If $a < b$, then $a + c < b + c$. This holds for $\leq, >, \geq$.
- **Multiplication ($c > 0$):** If $a < b$, then $ac < bc$. This holds for $\leq, >, \geq$.
- **Multiplication ($c < 0$):** If $a < b$, then $ac > bc$. This holds for $\leq, >, \geq$ (the sign flips).
- Subtraction follows the rules of addition. Division follows the rules of multiplication.

6 Summation and Product Notations

- **Summation Notation:** In $\sum_{i=0}^k s(i)$, i is the *index*, $i = 0$ is the *lower limit*, k is the *upper limit*, and $s(i)$ is the sequence we are summing.
- **Product Notation:** In $\prod_{i=0}^k s(i)$, everything is the same as summation, except we use π to indicate that we multiply the sequence.

7 Integer Division

- **Modulo** - Denoted by % or mod , the modulus operator gives the remainder of an integer division. E.g. $10 \% 4 = 2$
- **Congruency** - a is congruent to b modulo m (denoted $a \equiv b \pmod{m}$), if $a \% m = b \% m$ or $(a - b) \% m = 0$
- **Divides:** - The "divides" operator, denoted $a|b$, returns **True** if $b \% a = 0$ and **False** otherwise.

8 Evens and Odds

- **Even** - An integer, n is even if there exists an integer k such that $n = 2k$ (or $2|n$, $n \% 2 = 0$, $n \equiv 0 \pmod{2}$)
- **Odd** - An integer, n is odd if there exists an integer k such that $n = 2k + 1$ (or $2 \nmid n$, $n \% 2 = 1$, $n \equiv 1 \pmod{2}$)

9 Logarithms and Exponents

Laws of Exponents and Logarithms:

- | | | |
|--|--|--------------------------------|
| (a) $w^{x+y} = w^x w^y$ | (b) $(w^x)^y = w^{xy}$ | (c) $v^x w^x = (vw)^x$ |
| (d) $\frac{w^x}{w^y} = w^{x-y}$ | (e) $\frac{v^x}{w^x} = \left(\frac{v}{w}\right)^x$ | (f) $\log_b(x^y) = y \log_b x$ |
| (g) $\log_b(xy) = \log_b x + \log_b y$ | (h) $\log_b\left(\frac{x}{y}\right) = \log_b x - \log_b y$ | (i) $b^{\log_b x} = x$ |
| (j) $\log_a x = \frac{\log_b x}{\log_b a}$ | (k) If $b^y = x$, then $\log_b x = y$ | |

10 Quadratic Equations

- **Quadratic Equation:** Equation of the form $ax^2 + bx + c$ where $a \neq 0$
- **Factoring Quadratics:** $(fx + d)(gx + e) = (fg)x^2 + (gd + fe)x + de$
- **Quadratic Formula:** $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

11 Number Systems

- Binary: Base 2, Digits 0,1
- Decimal: Base 10, Digits 0-9
- Octal: Base 8, Digits 0-7
- Hexadecimal: Base 16, Digits 0-9,A-F