
Proof Examples

Example 1

Conjecture: Every odd integer is the difference of two squares.

Proof (Direct): Let n be an odd integer. $\exists_{k \in \mathbb{Z}}$ s.t. $n = 2k + 1$

To gain some insight:

$$3 = 2(1) + 1 = 2^2 - 1^2, \quad 5 = 2(2) + 1 = 3^2 - 2^2,$$

$$7 = 2(3) + 1 = 4^2 - 3^2, \quad 27 = 2(13) + 1 = 14^2 - 13^2$$

Observation 1: odd numbers seem to be the difference of two consecutive squares.

Observation 2: For an odd number, $n = 2k + 1$, it seems to be the sum of the squares $(k + 1)^2 - k^2$.

Example 1

Conjecture: Every odd integer is the difference of two squares.

Proof (Direct): Let n be an odd integer. $\exists_{k \in \mathbb{Z}}$ s.t. $n = 2k + 1$

To gain some insight:

$$3 = 2(1) + 1 = 2^2 - 1^2, \quad 5 = 2(2) + 1 = 3^2 - 2^2,$$

WARNING: We have not proved this yet!

$$7 = 2(3) + 1 = 4^2 - 3^2, \quad 27 = 2(13) + 1 = 14^2 - 13^2$$

Observation 1: odd numbers seem to be the difference of two consecutive squares.

Observation 2: For an odd number, $n = 2k + 1$, it seems to be the sum of the squares $(k + 1)^2 - k^2$.

Example 1

Conjecture: Every odd integer is the difference of two squares.

Proof (Direct): Let n be an odd integer.

$$\exists_{k \in \mathbb{Z}} \text{ s.t. } n = 2k + 1$$

Observe: $(k + 1)^2 - k^2 = k^2 + 2k + 1 - k^2 = 2k + 1$

So, when we have $n = 2k + 1$, we will add and subtract k^2 to the right side: $n = 2k + 1 + k^2 - k^2$

Which can be factored to: $n = (k + 1)^2 - k^2$.

Therefore, every odd integer is the difference of two squares.

Example 2(a)

Conjecture: If x is irrational, then $\frac{1}{x}$ is irrational.

Proof (Contrapositive): Assume $\frac{1}{x}$ is rational.

By definition, $\exists_{p,q \in \mathbb{Z}}$ s.t. $\frac{1}{x} = \frac{p}{q}$

Solving for x , we get $x = \frac{q}{p}$. Since p and q are both integers, x is rational.

Thus we have shown that the contrapositive is true.

Therefore, if x is irrational, then $\frac{1}{x}$ is irrational.

Example 2(b)

Conjecture: If x is irrational, then $\frac{1}{x}$ is irrational.

Proof (Contradiction): Assume x is irrational but $\frac{1}{x}$ is rational.

By definition, $\exists_{p,q \in \mathbb{Z}}$ s.t. $\frac{1}{x} = \frac{p}{q}$.

Solving for x we get $x = \frac{p}{q}$. We know p and q are integers so this makes x rational, which is a contradiction.

Therefore, If x is irrational, then $\frac{1}{x}$ is irrational.

Example 3

Conjecture: Pick a list of 22 days in a year. At least four of those days fall on the same day of the week.

Proof (Contradiction): Assume not. Assume that in our list of 22 days that no day of the week occurs more than 3 times.

Let d_i be the number of times i^{th} day of the week occurs, where $1 \leq i \leq 7$.

If $d_i \leq 3$, then $\sum_{i=1}^7 d_i \leq \sum_{i=1}^7 3 \leq 7 * 3 = 21$. This is a

contradiction because we assumed our list had 22 days on it.

Therefore, if we pick a list of 22 days in a year, then at least four of those days fall on the same day of the week.

Example 4

Conjecture: If $x, y \in \mathbb{R}$, then $|x| + |y| \geq |x + y|$

Proof (By Cases): Assume x and y are real numbers.

We know that if $x \geq 0$, $|x| = x$ and if
 $x < 0$, $|x| = -x$

Case 1: $x \geq 0, y \geq 0$.

$$|x| + |y| = x + y.$$

$$|x + y| = x + y$$

Thus $|x| + |y| \geq |x + y|$

Example 4

Conjecture: If $x, y \in \mathbb{R}$, then $|x| + |y| \geq |x + y|$

Proof (By Cases): Assume x and y are real numbers.

We know that if $x \geq 0$, $|x| = x$ and if $x < 0$, $|x| = -x$

Case 2: $x \geq 0$, $y < 0$, $x + y \geq 0$.

$$|x| + |y| = x + (-y) = x - y.$$

$-y \geq y$, so $x + -y \geq x + y$.

Since $x + y \geq 0$, $|x + y| = x + y$.

Thus $|x| + |y| = x + -y \geq x + y = |x + y|$

Example 5

Conjecture: If $x, y \in \mathbb{R}$, then $|x| + |y| \geq |x + y|$

Proof (By Cases): Assume x and y are real numbers.

We know that if $x \geq 0$, $|x| = x$ and if $x < 0$, $|x| = -x$

Case 3: $x \geq 0$, $y < 0$, $x + y < 0$.

$$|x| + |y| = x + (-y) = x - y.$$

Since $x + y < 0$, $|x + y| = -(x + y) = -x - y$.

$$x \geq -x \text{ so } x - y \geq -x - y$$

$$\text{Thus } |x| + |y| = x - y \geq -x - y = |x + y|$$

Example 5

Conjecture: If $x, y \in \mathbb{R}$, then $|x| + |y| \geq |x + y|$

Proof (By Cases): Assume x and y are real numbers.

Case 4 & 5: Same as case 2 & 3 with x and y flipped

Case 6: $x < 0, y < 0$.

$$|x| + |y| = -x - y.$$

$$|x + y| = -(x + y) = -x - y.$$

Thus, $|x| + |y| \geq |x + y|$

Therefore, $|x| + |y| \geq |x + y|$ in all cases.

Example 6

Conjecture: The following three statements about $x \in \mathbb{R}$ are equivalent: (i) x is rational, (ii) $x/2$ is rational, and (iii) $3x - 1$ is rational.

Proof: To show that these three are equivalent, it is sufficient to show (i) \rightarrow (ii), (ii) \rightarrow (iii), and (iii) \rightarrow (i).

Why is that sufficient?

1. $(i) \rightarrow (iii) \equiv (i) \rightarrow (ii) \wedge (ii) \rightarrow (iii)$
2. $(ii) \rightarrow (i) \equiv (ii) \rightarrow (iii) \wedge (iii) \rightarrow (i)$
3. $(iii) \rightarrow (ii) \equiv (iii) \rightarrow (i) \wedge (i) \rightarrow (ii)$

Example

Conjecture: The following three statements about $x \in \mathbb{R}$ are equivalent: (i) x is rational, (ii) $x/2$ is rational, and (iii) $3x - 1$ is rational.

Proof: (i) \rightarrow (ii) (direct): Assume x is rational.

By definition, $\exists_{p,q \in \mathbb{Z}}$ s.t. $x = \frac{p}{q}$.

$x/2 = \frac{\frac{p}{q}}{2} = \frac{p}{2q}$. $2q$ is an integer, therefore $x/2$ is rational.

Therefore, (i) \rightarrow (ii).

(continued)

Example

Conjecture: The following three statements about $x \in \mathbb{R}$ are equivalent: (i) x is rational, (ii) $x/2$ is rational, and (iii) $3x - 1$ is rational.

Proof: (ii) \rightarrow (iii) (direct): Assume $x/2$ is rational.

By definition, $\exists_{p,q \in \mathbb{Z}}$ s.t. $x/2 = \frac{p}{q}$.

$$x = \frac{2p}{q}, \quad 3x - 1 = 3\left(\frac{2p}{q}\right) - 1 = \frac{6p}{q} - 1 = \frac{6p}{q} - \frac{q}{q} = \frac{6p - q}{q}$$

$6p - q \in \mathbb{Z}$, so $3x - 1$ is rational.

Therefore, (ii) \rightarrow (iii).

(continued)

Example

Conjecture: The following three statements about $x \in \mathbb{R}$ are equivalent: (i) x is rational, (ii) $x/2$ is rational, and (iii) $3x - 1$ is rational.

Proof: (iii) \rightarrow (i) (direct): Assume $3x - 1$ is rational.

By definition, $\exists_{p,q \in \mathbb{Z}}$ s.t. $3x - 1 = \frac{p}{q}$.

$$3x = \frac{p}{q} + 1 = \frac{p + q}{q}. \quad x = \frac{\frac{p + q}{q}}{3} = \frac{p + q}{3q}.$$

$p + q \in \mathbb{Z}$, $3q \in \mathbb{Z}$ so x is rational.

Therefore, (iii) \rightarrow (i).

Thus, the statements x is rational, $x/2$ is rational, and $3x - 1$ is rational are equivalent.