
Sequences and Strings

4.3

Sequences

Definition: Sequence [1st Attempt]

An ordered list of items

Notation:

- Labels are lower-case letters
- Elements are subscripted: e_1, e_2, \dots
- $\{e_n\} \Rightarrow e$ is an n -element sequence.

Example: Soup!

Cost sequence: $s = 2, 4, 6, 8, 10, \dots$ (\$2 per can)

Soup Saturday: Buy 3 cans of soup, get one free!

$s' = 2, 4, 6, 6, 8, 10, 12, 12, \dots$ (Not a set!)

Rules

Recall: $\sum_{i=1}^n 2i$ ← Sequence defined by the rule $2i$

Example:

$s_n = 2n$ defines the original soup price sequence

$n^2 + 1, n \geq 0$ defines the infinite sequence

1, 2, 5, 10, 17, ...

More notation:

Infinite sequences:

1. Ellipses (as in 1, 2, 5, 10, 17, ...)

2. $\{d_n\}_{n=1}^{\infty}$

Sequences and Functions

Definition: Sequence [Final Version]

A sequence is the ordered range of a function from a set of integers to some set S

Example:

$o(n) = 2n - 1$ on the domain $\{1,2,3,4,5\}$ defines the sequence 1,3,5,7,9

As a relation: $\{(1,1), (2,3), (3,5), (4,7), (5,9)\}$

Range of $\{1,3,5,7,9\}$

(Thus, the “ordered range” wording)

Arithmetic and Geometric Sequences

Definition: Arithmetic Sequence (a.k.a. Arithmetic Progression)

In an arithmetic sequence, the *common difference* $d = a_{n+1} - a_n$ is constant

Definition: Geometric Sequence (a.k.a. Geometric Progression)

In a geometric sequence, the *common ratio* $r = \frac{g_{n+1}}{g_n}$ is constant

Example:

In a : 1, 3, 5, 7, 9 $d = 2$

In g : $1, \frac{2}{3}, \frac{4}{9}, \frac{8}{27}, \dots$, $r = \frac{2}{3}$

Arithmetic Series

- The sum of the terms of an arithmetic sequence (a.k.a arithmetic series):

$$s_n = a_1 + \dots + a_n = \frac{1}{2}n(a_1 + a_n)$$

- Here's why: First, note that $a_n = a_1 + (n - 1)d$.

- Next, here are two expressions for s_n :

- $s_n = a_1 + (a_1 + d) + (a_1 + 2d) + \dots + (a_1 + (n - 1)d)$

- $s_n = (a_n - (n - 1)d) + (a_n - (n - 2)d) + \dots + (a_n - d) + a_n$

- Sum these expressions, and the d terms cancel, leaving:

- $2s_n = na_1 + na_n$ or $s_n = \frac{1}{2}n(a_1 + a_n)$

- Ex: In 1,3,5 $d = 2$, $a_1 = 1$, $a_1 + d = 3$ and $a_1 + 2d = 5$

Increasing Sequences

Definition: Increasing Sequence

An increasing sequence labeled i is ordered such that $i_n \leq i_{n+1}$.

Definition: Non-Decreasing Sequence

A non-decreasing sequence labeled i is ordered such that $i_n \leq i_{n+1}$ [Same as increasing!]

Definition: Strictly Increasing Sequence

A strictly increasing sequence labeled i is ordered such that $i_n < i_{n+1}$

Decreasing Sequences

Definition: Decreasing Sequence

A decreasing sequence labeled i is ordered such that $i_n \geq i_{n+1}$.

Definition: Non-Increasing Sequence

A non-increasing sequence labeled i is ordered such that $i_n \geq i_{n+1}$ [Same as decreasing!]

Definition: Strictly Decreasing Sequence

A strictly decreasing sequence labeled i is ordered such that $i_n > i_{n+1}$

Examples: Increasing/Decreasing Sequences

- The sequence $g = 1, 2, 2, 2, 6, 8, 8, 9$ is:
 - Increasing
 - Non-Decreasing
-

- $h_n = \frac{1}{n}, 4 \leq n \leq 7$ ($h = \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \frac{1}{7}$)

- $\{h_n\}_{n=4}^7$ is:
 - Decreasing, Non-Increasing
 - Strictly Decreasing

Subsequences

Definition: Subsequence

Sequence x is a subsequence of sequence y when the elements of x are found within y in the same relative order

Example:

Is $\frac{1}{4}, \frac{1}{6}$ a subsequence of $\{h_n\}_{n=4}^7$?

Is 8,6,2 a subsequence of $g = 1,2,2,2,6,8,8,9$?

Need to Identify a Sequence?

A great resource for sequences:

The Online Encyclopedia of Integer Sequences

(<http://oeis.org/>)

Example:

Let's try it! 2,3,5,7,11,13,17

Strings

Definition: *String*

A string is a contiguous finite sequence of 0 or more elements drawn from a set called the *alphabet*

Example:

A sequence of DNA nucleotides (e.g. ATTGACCT) is called a string.

A Java String also qualifies (alphabet: UNICODE values)

Strings

- Notation:
 - Lambda (λ) represents the empty (null) string
 - xy means strings x and y are concatenated
 - Superscripts denote repetition of concatenation
 - $|x|$ represents the length of string x
 - A^* is the set of strings that can be formed using elements of an alphabet A
 - A^* is an infinite set
 - $\lambda \in A^*$

Set Cardinality Revisited

An observation about set cardinality:

Two sets A and B have the same cardinality **iff** there is a bijection from A to B

Definition: Finite

A set S is finite if there exists a bijective mapping between it and a set of cardinality $|S|$

Set Cardinality Revisited

Definition: Countably Infinite (a.k.a. Denumerably Infinite)

A set is countably infinite if there exists a bijective mapping between the set and either \mathbb{Z}^* or \mathbb{Z}^+

Definition: Countable

A set is countable if it is finite or countably infinite.
(Otherwise, it is *uncountable*.)

Set Cardinality Revisited

Example:

Is the set of digits in the 'house number' of the Gould-Simpson building countable?

$$1040 \text{ E. 4th St.} \Rightarrow \{0,1,4\}$$

This set is finite, so, yes!


Is the set of positive multiples of 5 countable?

$$\begin{array}{cccccccc} \mathbb{Z}^+ : & 1 & 2 & 3 & 4 & \dots & 5z & y \\ & & & & & & & \frac{y}{5} \\ & 5 & 10 & 15 & 20 & \dots & z & \end{array}$$

Invertible \rightarrow Bijection \rightarrow Countable! (So, yes!)

Set Cardinality Revisited

Question: Are the positive rational numbers countable?



Just skip duplicates

$$R = \left\{ \left(1, \frac{1}{1}\right), \left(2, \frac{2}{1}\right), \left(3, \frac{1}{2}\right), \left(4, \frac{1}{3}\right), \left(5, \frac{3}{1}\right), \dots \right\}$$

$$R^{-1} = \left\{ \left(\frac{1}{1}, 1\right), \left(\frac{2}{1}, 2\right), \left(\frac{1}{2}, 3\right), \left(\frac{1}{3}, 4\right), \left(\frac{3}{1}, 5\right), \dots \right\}$$

It is invertible, therefore it is a bijection.

Yes: (This is an application of a 'pairing function which invertible maps after duplicates are removed.) (With duplicates, the function is not invertible!)

(This is an example of a boustrophedonic path.)

Set Cardinality Revisited

Conjecture: A pairing function for \mathbb{R} cannot exist

Proof (Contradiction): Assume that a pairing function for the reals does exist. Given the set of real numbers, form a new number (not in the pairing) by changing the d^{th} digit of the d^{th} number. For example:

0	.	8	9	4	6	...
0	.	1	2	0	4	...
0	.	7	7	7	6	...
0	.	4	5	6	7	...
⋮	.	⋮	⋮	⋮	⋮	⋮
0	.	5	4	9	1	...

**0.5491... won't be in pairing,
because it has 1 digit
different than every number
in the pairing**

The result must be a value not in the set of reals, yet it is a real. This is a contradiction (This is called *Cantor's Diagonal Argument*).

Therefore, a pairing function for \mathbb{R} cannot exist